

The tropical critical points of an affine matroid

Turn panic into magic

SIAM AG 2023, Eindhoven University of Technology, Eindhoven

Raul Penagiao

MPI MiS Leipzig

July 10th, 2023

Slides can be found at raulpenagiao.github.io/
Joint work with Federico Ardila and Chris Eur

Optimization of a monomial

Fix some vector $\mathbf{w} = (w_1, \dots, w_n) \in \mathbb{R}^n$.

Optimize $f_{\mathbf{w}} : \mathbf{x} \mapsto x_1^{w_1} \cdots x_n^{w_n}$ on a variety $X \subset (\mathbb{C}^*)^n$.




Figure: A variety where we can optimize $f_{\mathbf{w}}$

What is the number of critical points of f ? Does it depend on the choice of \mathbf{w} ? For generic \mathbf{w} , no!

This number is called the **maximum likelihood degree** of a model X .
If X is a vector space, $\text{MLDeg}(X) = \beta(M(V))$.

Edge weight problem

Given $G = (V, E)$, fix some vector $\mathbf{w} = (w_1, \dots, w_n) \in \mathbb{R}^n$.




Figure: Find x and y edge weights that are *compatible* with G and $(G \setminus z)^*$.

- The sum of the weights is \mathbf{w} .
- **(Compatible)** Every cycle has at least two minimal edges.

Fix $\mathbf{w} = (0, 1, 1, 2, 2, 5, 3, 4, 7)$.





Figure: Find \mathbf{x} and \mathbf{y} edge weights that are *compatible* with G and $(G \setminus z)^*$.

	a	b	c	d	e	f	g	h	i
\mathbf{x}	00	01	00	00	02	03	00	00	02
\mathbf{y}	000	110	111	222	20	52	33	44	75
\mathbf{w}	0	1	1	2	2	5	3	4	7

- 1 Introduction
- 2 Matroids
- 3 The Bergman Fan
- 4 Degree of Bergman Fan

Graphical matroid

Given a graph $G = (V, E)$, the collection of edges E forms a matroid.



Independent sets \mapsto forests

Basis \mapsto Spanning forests


Circuits \mapsto Simple cycles

Rank of set $A \subseteq E$ \mapsto size of largest spanning forest

Flats

Maximal sets with a fixed rank.

That is, F is a flat if for any $i \notin F$, $r_M(F \cup i) > r_M(F)$.



\emptyset , matchings , complete subgraphs , . . .

$$\{\emptyset \subsetneq a \subsetneq za \subsetneq zafg \subsetneq zabcdefghi\}$$

The uniform matroid

Basis of the uniform matroid $U_{n,k}$ = all sets of size k in $[n]$.
 Any set of size $\leq k$ is independent.




Figure: Matroid $U_{7,5}$ along with a basis B and independent set I .

Any set of size $\leq k - 1$ is a flat.

Any complete flag of flats is of the form

$$\{\emptyset \subsetneq \{v_1\} \subsetneq \{v_1, v_2\} \subsetneq \cdots \subsetneq \{v_1, \dots, v_{k-1}\} \subsetneq [n]\}$$

The Bergman Fan

$\Sigma_M := \{\vec{x} \in \mathbb{R}^n \mid \forall \text{ circuits } C \text{ we have } \min_{c \in C} x_c \text{ is attained twice}\}.$

$\Sigma_M := \{\vec{x} \in \mathbb{R}^n / \mathbb{1}_{\mathbb{R}} \mid \forall \text{ circuits } C \text{ we have } \min_{c \in C} x_c \text{ is attained twice}\}.$

If $M = U_{n,k}$, any set of size $k+1$ is a circuit.

A point $\vec{x} \in \mathbb{R}^n$ is on the Bergman fan if $\#\{i \in E \mid x_i > \min \vec{x}\} \leq k-1$.

$$\Sigma_M = \bigcup_{|I|=n-k+1} \{\vec{x} \in \mathbb{R}^n / \mathbb{1}_{\mathbb{R}} \mid \arg \min \vec{x} \subseteq I\}.$$

$$\Sigma_{U_{3,2}} = \{(a, a, b) \mid a \leq b\} \cup \{(a, b, a) \mid a \leq b\} \cup \{(b, a, a) \mid a \leq b\}.$$




Figure: Two elements in the Bergman fan of the graphical matroid

Theorem (Sturmfels and Feichtner, 2004)

The Bergman Fan of a matroid decomposes into the following cones

$$\Sigma_M = \bigcup_{\mathcal{F} \text{ flag of flats}} \mathcal{C}_{\mathcal{F}} = \bigcup_{\substack{F_1 \subset \dots \subset F_k \\ \text{flag of flats}}} \{x_i \geq x_j \text{ whenever } i \in F_k, j \notin F_k\}.$$

Problem

Can we compute the degree of the tropical variety Σ_M ?




Figure: A variety X : Degree = # of intersections with a **line** in \mathbb{C}^n .

Just intersect it with a hyperplane with dimension = $n - \dim X$

Note: Hyperplanes in the tropical world are Bergman fans of $U_{n,k}$.

Example of degree computation

Consider M the graphical matroid of K_5 , of rank 4 (so $\dim \Sigma_M = 3$). One has $\Sigma_M \subseteq \mathbb{R}^{10}/\mathbb{R}\mathbf{1}$, we intersect it with a hyperplane of $\dim = 6$.

$$|\Sigma_{U_{10,7}} + \underbrace{(1, 10, 100, 1000, \dots, 10^9)}_{\vec{\omega}} \cap \Sigma_M| = ?$$




Figure: There is only one $x \in \Sigma_M$ and $y \in \Sigma_U$ such that $y + w = x$. Such vector x belongs to a **cone determined by the greedy basis of M** .

Activities

Fix total order in V , ground set of a matroid M , basis B .

- $e \in B$ is internal activity if $e = \min C^\perp$, where $C^\perp \subseteq B^c \cup e$ is a cocircuit (a cut of the graph).
- $e \notin B$ is external activity if $e = \min C$, where $C \subseteq B \cup e$ is a circuit.




Figure: Fix order $z < a < b < c < d < e < f < g < h < i$.

$i(befg) = 0$, $e(befg) = 2$, $i(zadf) = 2$, $e(zadf) = 1$

$i(B) = 0$ nbc basis. $i(B) = 0$ and $e(B) = 1$ β -nbc basis

Degree computations

Theorem (Greedy basis algorithm)

$$\deg(\Sigma_M) = \Sigma_M \cap (\mathbf{w} + \Sigma_{U_{n,n-k-1}}) = 1$$

Theorem (Adiprasito, Huh, Katz, 2018)

$$\deg(-\Sigma_M) = (-\Sigma_M) \cap (\mathbf{w} + \Sigma_{U_{n,n-k-1}}) = \#\{\text{nbc bases}\}$$

Theorem (Agostini, Brysiewicz, Fevola, Kühne, Sturmfels, Telen 2021 and Ardila, Eur, P 2022)



$$\deg(\Sigma_{(M/0)^\perp} \cdot -\Sigma_M) = (-\Sigma_M) \cap (\mathbf{w} + \Sigma_{(M/0)^\perp}) = \#\{\beta - \text{nbc bases}\}$$

Finding Nemo points from nbc bases

$$\deg(\Sigma_{(M/0)^\perp} \cdot -\Sigma_M) = ?$$

Find $\mathbf{x} \in \Sigma_M$, $\mathbf{y} \in \Sigma_{(M/0)^\perp}$ such that $\mathbf{x} + \mathbf{y} = \mathbf{w}$.

Fix generic $\mathbf{w} = (10^{i-1})_i$.



Find a β -nbc basis $B = zbeg \rightarrow g|e|bd|zacfhi$,

corresponding β -nbc cobasis $B^\perp = acdfhi \rightarrow i|h|f|gd|c|eba$.

	a	b	c	d	e	f	g	h	i
\mathbf{x}	0	9	0	9	$10^4 - 1$	0	$10^6 - 10^3 + 9$	0	0
\mathbf{y}	1	1	100	$10^3 - 9$	1	10^5	$10^3 - 9$	10^7	10^8
\mathbf{w}	1	10	100	10^3	10^4	10^5	10^6	10^7	10^8

Biblio

- Agostini D., Brysiewicz T., Fevola C., Kühne L., Sturmfels B., Telen S. (2021). *Likelihood Degenerations*. Motivation behind ML degree computations
- Adiprasito K., Huh J., Katz E. (2018). *Hodge Theory for combinatorial geometries* Degree computations in matroids
- Ardila F., Eur C., RP (2022) *The maximum likelihood of a matroid.*

Thank you

